



News, information, and educational materials about the world's synchrotron and free electron laser **light source facilities**

SEARCH

LEARN ABOUT LIGHT SOURCES

LIGHT SOURCE NEWS

LIGHT SOURCE SCIENCE

FOR LIGHT SOURCE USERS

Printer Friendly Version

What Is a Light Source?

News Archive

Press Releases

Light Sources in the News

News by Date

News by Subject

News by Facility

Information about Light Sources

Image Bank

Educational Material

Facility Information

Useful Links

Conferences etc.

Courses

### Subscribe to NEWS FLASH

Subscribe now to keep up to date with breaking news from the light source community.

RSS

**-Press Release Number:** PR-NSF-07-1  
**-Source:** [National Science Foundation](#)  
**-Date issued:** January 8, 2007  
**-Contact:** Cheryl Dybas, NSF +1 (703) 292-7734, [cdybas@nsf.gov](mailto:cdybas@nsf.gov)

## Diamonds from Outer Space: Geologists Discover Origin of Earth's Mysterious Black Diamonds

If indeed "a diamond is forever," the most primitive origins of Earth's so-called black diamonds were in deep, universal time, geologists have discovered. Black diamonds came from none other than interstellar space.

In a paper published online on December 20, 2006, in the journal *Astrophysical Journal Letters*, scientists Jozsef Garai and Stephen Haggerty of Florida International University, along with Case Western Reserve University researchers Sandeep Rekhi and Mark Chance, claim an extraterrestrial origin for the unique black diamonds, also called carbonado diamonds.

Infrared synchrotron radiation at Brookhaven National Laboratory was used to discover the diamonds' source.

"Trace elements critical to an 'ET' origin are nitrogen and hydrogen," said Haggerty. The presence of hydrogen in the carbonado diamonds indicates an origin in a hydrogen-rich interstellar space, he and colleagues believe.

**Image 1.** Black, or carbonado, diamonds, came from outer space, geologists have discovered. (Photo: Steve Haggerty)



The term carbonado was coined by the Portuguese in Brazil in the mid-18th century; it's derived from its visual similarity to porous charcoal. Black diamonds are found only in Brazil and the Central African Republic.

"Conventional diamonds are mined from explosive volcanic rocks [kimberlites] that transport them from depths in excess of 100 kilometers to the Earth's surface in a very short amount of time," said Sonia Esperanca, program director in the National Science Foundation's Division of Earth Sciences, which funded the research. "This process preserves the unique crystal structure that makes diamonds the hardest natural material known."

From Australia to Siberia, from China to India, the geological settings of conventional diamonds are virtually identical, said Haggerty. None of them are compatible with the formation of black diamonds.

Approximately 600 tons of conventional diamonds have been mined, traded, polished and adorned since 1900. "But not a single black/carbonado diamond has been discovered in the world's mining fields," Haggerty said.

The new data support earlier research by Haggerty showing that carbonado diamonds formed in stellar supernovae explosions. Black diamonds were once the size of asteroids, a kilometer or more in diameter when they first landed on Earth.

###

*The National Science Foundation (NSF) is an independent federal agency that supports fundamental research and education across all fields of science and engineering, with an annual budget of \$5.58 billion. NSF funds reach all 50 states through grants to nearly 1,700 universities and institutions. Each year, NSF receives about 40,000 competitive requests for funding, and makes nearly 10,000 new funding awards. The NSF also awards over \$400 million in professional and service contracts yearly.*

The original press release can be found at: [http://www.nsf.gov/news/news\\_summ.jsp?cntn\\_id=108270](http://www.nsf.gov/news/news_summ.jsp?cntn_id=108270)

[**Note:** Published in *Astrophysical Journal Letters*, 2006 December 20, Volume 653:L153-L156, Number 2, Part 2, DOI: [10.1086/510451](https://doi.org/10.1086/510451); Jozsef Garai, Stephen E. Haggerty, Sandeep Rekhi, and Mark Chance ; "Infrared Absorption Investigations Confirm the Extraterrestrial Origin of Carbonado Diamonds"

The synchrotron IR experiment was carried out at the U2B beamline at the [National Synchrotron Light Source](#) (Brookhaven National Laboratory). Brookhaven National Laboratory is funded by the U.S. Department of Energy's Office of Science.

*The contents of all materials on lightsources.org are the sole responsibility of the authors of the materials and/or the facilities or institutions under whose auspices the materials were produced. -lightsources.org]*